



Enumeration of Domino Tilings of an Aztec Rectangle with boundary defects

arXiv:1605.09169

Manjil Saikia (Universität Wien)

01 August, 2016

3rd AEC Summer School 2016, RISC, Hagenberg

Table of contents

1. Introduction and Motivation
2. Graphical Condensation
3. Aztec Rectangles with Defects

Introduction and Motivation

Aztec Diamonds

- In 1991, Elkies, Kuperberg, Larsen and Propp introduced a new class of object which they called Aztec Diamonds.

Aztec Diamonds

- In 1991, Elkies, Kuperberg, Larsen and Propp introduced a new class of object which they called Aztec Diamonds.
- The Aztec Diamond of order n (denoted by $AD(n)$) is the union of all unit squares inside the contour $|x| + |y| = n + 1$

Aztec Diamonds

- In 1991, Elkies, Kuperberg, Larsen and Propp introduced a new class of object which they called Aztec Diamonds.
- The Aztec Diamond of order n (denoted by $AD(n)$) is the union of all unit squares inside the contour $|x| + |y| = n + 1$




Figure 1: $AD(3)$, Aztec Diamond of order 3

Aztec Diamond Theorem

- A domino is the union of any two unit squares sharing an edge, and a domino tiling of a region is a covering of the region by dominoes so that there are no gaps or overlaps.

Aztec Diamond Theorem

- A domino is the union of any two unit squares sharing an edge, and a domino tiling of a region is a covering of the region by dominoes so that there are no gaps or overlaps.
- They considered the problem of counting the number of domino tiling the Aztec Diamond with dominoes and presented four different proofs of the following result.

Aztec Diamond Theorem

- A domino is the union of any two unit squares sharing an edge, and a domino tiling of a region is a covering of the region by dominoes so that there are no gaps or overlaps.
- They considered the problem of counting the number of domino tiling the Aztec Diamond with dominoes and presented four different proofs of the following result.

Theorem (Elkies–Kuperberg–Larsen–Propp)

The number of domino tilings of an Aztec Diamond of order n is $2^{n(n+1)/2}$.

Aztec Rectangles

- We denote by $\mathcal{AR}_{a,b}$ the Aztec rectangle which has a unit squares on the southwestern side and b unit squares on the northwestern side.

Aztec Rectangles

- We denote by $\mathcal{AR}_{a,b}$ the Aztec rectangle which has a unit squares on the southwestern side and b unit squares on the northwestern side.
- We assume $b \geq a$ unless otherwise mentioned. For $a < b$, $\mathcal{AR}_{a,b}$ does not have any tiling by dominoes.

Aztec Rectangles

- We denote by $\mathcal{AR}_{a,b}$ the Aztec rectangle which has a unit squares on the southwestern side and b unit squares on the northwestern side.
- We assume $b \geq a$ unless otherwise mentioned. For $a < b$, $\mathcal{AR}_{a,b}$ does not have any tiling by dominoes.
- The non-tileability of the region $\mathcal{AR}_{a,b}$ becomes evident if we look at the checkerboard representation of $\mathcal{AR}_{a,b}$

Aztec Rectangle




Figure 2: Checkerboard representation of an Aztec Rectangle with $a = 4, b = 10$

Aztec Rectangle Theorem

If we remove $b - a$ unit squares from the southeastern side then we have a simple product formula found by Mills, Robbins and Rumsey.

Aztec Rectangle Theorem

If we remove $b - a$ unit squares from the southeastern side then we have a simple product formula found by Mills, Robbins and Rumsey.

Theorem (Mills-Robbins-Rumsey, Elkies-Kuperberg-Larson-Propp)

Let $a < b$ be positive integers and $1 \leq s_1 < s_2 < \dots < s_a \leq b$. Then the number of domino tilings of $\mathcal{AR}_{a,b}$ where all unit squares from the south-eastern side are removed except for those in positions s_1, s_2, \dots, s_a is

$$2^{a(a+1)/2} \prod_{1 \leq i < j \leq a} \frac{s_j - s_i}{j - i}.$$

Aztec Rectangle Theorem

If we remove $b - a$ unit squares from the southeastern side then we have a simple product formula found by Mills, Robbins and Rumsey.

Theorem (Mills-Robbins-Rumsey, Elkies-Kuperberg-Larson-Propp)

Let $a < b$ be positive integers and $1 \leq s_1 < s_2 < \dots < s_a \leq b$. Then the number of domino tilings of $\mathcal{AR}_{a,b}$ where all unit squares from the south-eastern side are removed except for those in positions s_1, s_2, \dots, s_a is

$$2^{a(a+1)/2} \prod_{1 \leq i < j \leq a} \frac{s_j - s_i}{j - i}.$$

Our goal here is to extend this result.

Graphical Condensation

For a graph G , $M(G)$ denotes the number of perfect matchings of G .

For a graph G , $M(G)$ denotes the number of perfect matchings of G .

Theorem (Eric Kuo)

Let G be a planar graph with four vertices w, x, y, z that appear in that cyclic order on a face of G .

For a graph G , $M(G)$ denotes the number of perfect matchings of G .

Theorem (Eric Kuo)

Let G be a planar graph with four vertices w, x, y, z that appear in that cyclic order on a face of G . Then

$$\begin{aligned} M(G) &= M(G - \{w, x, y, z\}) + M(G - \{w, y\}) M(G - \{x, z\}) \\ &= M(G - \{w, x\}) M(G - \{y, z\}) + M(G - \{w, z\}) M(G - \{x, y\}). \end{aligned}$$

If $G = (V_1, V_2, E)$ is bipartite, and

If $G = (V_1, V_2, E)$ is bipartite, and

- $w, y \in V_1, x, z \in V_2, |V_1| = |V_2|$; second term vanishes

Theorem (Eric Kuo)

Let G be a planar graph with four vertices w, x, y, z that appear in that cyclic order on a face of G . Then

$$\begin{aligned} & M(G) M(G - \{w, x, y, z\}) + M(G - \{w, y\}) M(G - \{x, z\}) \\ &= M(G - \{w, x\}) M(G - \{y, z\}) + M(G - \{w, z\}) M(G - \{x, y\}). \end{aligned}$$

If $G = (V_1, V_2, E)$ is bipartite, and

- $w, y \in V_1, x, z \in V_2, |V_1| = |V_2|$; second term vanishes

If $G = (V_1, V_2, E)$ is bipartite, and

- $w, y \in V_1, x, z \in V_2, |V_1| = |V_2|$; second term vanishes
- $w, x \in V_1, y, z \in V_2, |V_1| = |V_2|$; third term vanishes

Theorem (Eric Kuo)

Let G be a planar graph with four vertices w, x, y, z that appear in that cyclic order on a face of G . Then

$$\begin{aligned} & M(G) M(G - \{w, x, y, z\}) + M(G - \{w, y\}) M(G - \{x, z\}) \\ &= M(G - \{w, x\}) M(G - \{y, z\}) + M(G - \{w, z\}) M(G - \{x, y\}). \end{aligned}$$

If $G = (V_1, V_2, E)$ is bipartite, and

- $w, y \in V_1, x, z \in V_2, |V_1| = |V_2|$; second term vanishes
- $w, x \in V_1, y, z \in V_2, |V_1| = |V_2|$; third term vanishes

If $G = (V_1, V_2, E)$ is bipartite, and

- $w, y \in V_1, x, z \in V_2, |V_1| = |V_2|$; second term vanishes
- $w, x \in V_1, y, z \in V_2, |V_1| = |V_2|$; third term vanishes
- $w, x, y, z \in V_1, |V_1| = |V_2| + 2$; first term vanishes

Theorem (Eric Kuo)

Let G be a planar graph with four vertices w, x, y, z that appear in that cyclic order on a face of G . Then

$$\begin{aligned} & M(G) M(G - \{w, x, y, z\}) + M(G - \{w, y\}) M(G - \{x, z\}) \\ &= M(G - \{w, x\}) M(G - \{y, z\}) + M(G - \{w, z\}) M(G - \{x, y\}). \end{aligned}$$

Theorem (Eric Kuo)

Let G be a planar graph with four vertices w, x, y, z that appear in that cyclic order on a face of G . Then

$$\begin{aligned} & M(G) M(G - \{w, x, y, z\}) + M(G - \{w, y\}) M(G - \{x, z\}) \\ &= M(G - \{w, x\}) M(G - \{y, z\}) + M(G - \{w, z\}) M(G - \{x, y\}). \end{aligned}$$

There is a generalization by Ciucu which we present.

Let $A = (a_{i,j})$ be a $2n \times 2n$ antisymmetric matrix and Γ_n be the set of all perfect matchings of K_{2n} .

Let $A = (a_{i,j})$ be a $2n \times 2n$ antisymmetric matrix and Γ_n be the set of all perfect matchings of K_{2n} . Then

$$\text{Pf}(A) = \sum_{\pi = \{(i_1, j_1), \dots, (i_n, j_n)\} \in \Gamma_n} \text{sgn } \pi \prod_{k=1}^n a_{i_k, j_k}$$

where $\text{sgn } \pi = \text{sgn } i_1 j_1 i_2 j_2 \dots i_n j_n$.

Let $A = (a_{i,j})$ be a $2n \times 2n$ antisymmetric matrix and Γ_n be the set of all perfect matchings of K_{2n} . Then

$$\text{Pf}(A) = \sum_{\pi = \{(i_1, j_1), \dots, (i_n, j_n)\} \in \Gamma_n} \text{sgn } \pi \prod_{k=1}^n a_{i_k, j_k}$$

where $\text{sgn } \pi = \text{sgn } i_1 j_1 i_2 j_2 \dots i_n j_n$.

- There are many ways to write π , so to see that $\text{Pf}(A)$ is well-defined we can assume that $i_k < j_k$ and $i_1 < i_2 < \dots < i_n$.

Let $A = (a_{i,j})$ be a $2n \times 2n$ antisymmetric matrix and Γ_n be the set of all perfect matchings of K_{2n} . Then

$$\text{Pf}(A) = \sum_{\pi = \{(i_1, j_1), \dots, (i_n, j_n)\} \in \Gamma_n} \text{sgn } \pi \prod_{k=1}^n a_{i_k, j_k}$$

where $\text{sgn } \pi = \text{sgn } i_1 j_1 i_2 j_2 \dots i_n j_n$.

- There are many ways to write π , so to see that $\text{Pf}(A)$ is well-defined we can assume that $i_k < j_k$ and $i_1 < i_2 < \dots < i_n$.
- Pfaffians have many interesting properties, such as

Let $A = (a_{i,j})$ be a $2n \times 2n$ antisymmetric matrix and Γ_n be the set of all perfect matchings of K_{2n} . Then

$$\text{Pf}(A) = \sum_{\pi = \{(i_1, j_1), \dots, (i_n, j_n)\} \in \Gamma_n} \text{sgn } \pi \prod_{k=1}^n a_{i_k, j_k}$$

where $\text{sgn } \pi = \text{sgn } i_1 j_1 i_2 j_2 \dots i_n j_n$.

- There are many ways to write π , so to see that $\text{Pf}(A)$ is well-defined we can assume that $i_k < j_k$ and $i_1 < i_2 < \dots < i_n$.
- Pfaffians have many interesting properties, such as

$$\text{Pf}(A)^2 = \det(A).$$

An Example

Let $n = 2$, then

An Example

Let $n = 2$, then

$$\text{Pf}(A) = \sum_{\pi = \{(i_1, j_1), (i_2, j_2)\} \in \Gamma_2} \text{sgn } \pi \prod_{k=1}^2 a_{i_k, j_k}$$

An Example

Let $n = 2$, then

$$\text{Pf}(A) = \sum_{\pi = \{(i_1, j_1), (i_2, j_2)\} \in \Gamma_2} \text{sgn } \pi \prod_{k=1}^2 a_{i_k, j_k}$$

$$\text{Pf}(A) = a_{1,2}a_{3,4} - a_{1,3}a_{2,4} + a_{1,4}a_{2,3}.$$

Theorem (Mihai Ciucu)

Let G be a planar graph with the vertices a_1, a_2, \dots, a_{2k} appearing in that cyclic order on a face of G .

Ciucu's Extension of Kuo's Condensation

Theorem (Mihai Ciucu)

Let G be a planar graph with the vertices a_1, a_2, \dots, a_{2k} appearing in that cyclic order on a face of G . Consider the skew-symmetric matrix $A = (a_{ij})_{1 \leq i, j \leq 2k}$ with entries given by

$$a_{ij} := M(G \setminus \{a_i, a_j\}), \text{ if } i < j. \quad (2.1)$$

Ciucu's Extension of Kuo's Condensation

Theorem (Mihai Ciucu)

Let G be a planar graph with the vertices a_1, a_2, \dots, a_{2k} appearing in that cyclic order on a face of G . Consider the skew-symmetric matrix $A = (a_{ij})_{1 \leq i, j \leq 2k}$ with entries given by

$$a_{ij} := M(G \setminus \{a_i, a_j\}), \text{ if } i < j. \quad (2.1)$$

Then we have that

$$M(G \setminus \{a_1, a_2, \dots, a_{2k}\}) = \frac{\text{Pf}(A)}{[M(G)]^{k-1}}. \quad (2.2)$$

Ciucu's Extension of Kuo's Condensation

Theorem (Mihai Ciucu)

Let G be a planar graph with the vertices a_1, a_2, \dots, a_{2k} appearing in that cyclic order on a face of G . Consider the skew-symmetric matrix $A = (a_{ij})_{1 \leq i, j \leq 2k}$ with entries given by

$$a_{ij} := M(G \setminus \{a_i, a_j\}), \text{ if } i < j. \quad (2.1)$$

Then we have that

$$M(G \setminus \{a_1, a_2, \dots, a_{2k}\}) = \frac{\text{Pf}(A)}{[M(G)]^{k-1}}. \quad (2.2)$$

We shall use this result in the next part.

Ciucu's Extension of Kuo's Condensation

Theorem (Mihai Ciucu)

Let G be a planar graph with the vertices a_1, a_2, \dots, a_{2k} appearing in that cyclic order on a face of G . Consider the skew-symmetric matrix $A = (a_{ij})_{1 \leq i, j \leq 2k}$ with entries given by

$$a_{ij} := M(G \setminus \{a_i, a_j\}), \text{ if } i < j. \quad (2.1)$$

Then we have that

$$M(G \setminus \{a_1, a_2, \dots, a_{2k}\}) = \frac{\text{Pf}(A)}{[M(G)]^{k-1}}. \quad (2.2)$$

We shall use this result in the next part.

A further common generalization of Kuo and Ciucu's results can be found in our preprint online.

Aztec Rectangles with Defects

Our Ideas

- We consider regions with defects (one unit square removed) on the boundaries of an Aztec Rectangle.

Our Ideas

- We consider regions with defects (one unit square removed) on the boundaries of an Aztec Rectangle.
- Mills-Robbins-Rumsey's result is for defects on one of the longer sides.

Our Ideas

- We consider regions with defects (one unit square removed) on the boundaries of an Aztec Rectangle.
- Mills-Robbins-Rumsey's result is for defects on one of the longer sides.
- It is well known, that domino tilings of Aztec Diamonds/Rectangles correspond to counting the number of perfect matchings of its dual planar graph.

Our Ideas

- We consider regions with defects (one unit square removed) on the boundaries of an Aztec Rectangle.
- Mills-Robbins-Rumsey's result is for defects on one of the longer sides.
- It is well known, that domino tilings of Aztec Diamonds/Rectangles correspond to counting the number of perfect matchings of its dual planar graph.
- So, from now on we use the terms matchings and tilings equivalently.

Our Ideas

- We consider regions with defects (one unit square removed) on the boundaries of an Aztec Rectangle.
- Mills-Robbins-Rumsey's result is for defects on one of the longer sides.
- It is well known, that domino tilings of Aztec Diamonds/Rectangles correspond to counting the number of perfect matchings of its dual planar graph.
- So, from now on we use the terms matchings and tilings equivalently.
- We would like to use Ciucu's result to give a Pfaffian for the regions we are interested in.

Our Ideas

- We consider regions with defects (one unit square removed) on the boundaries of an Aztec Rectangle.
- Mills-Robbins-Rumsey's result is for defects on one of the longer sides.
- It is well known, that domino tilings of Aztec Diamonds/Rectangles correspond to counting the number of perfect matchings of its dual planar graph.
- So, from now on we use the terms matchings and tilings equivalently.
- We would like to use Ciucu's result to give a Pfaffian for the regions we are interested in.
- But the problem is, if $k > 0$ in Ciucu's result, then our graph G has no matchings.

Our Ideas

- We consider regions with defects (one unit square removed) on the boundaries of an Aztec Rectangle.
- Mills-Robbins-Rumsey's result is for defects on one of the longer sides.
- It is well known, that domino tilings of Aztec Diamonds/Rectangles correspond to counting the number of perfect matchings of its dual planar graph.
- So, from now on we use the terms matchings and tilings equivalently.
- We would like to use Ciucu's result to give a Pfaffian for the regions we are interested in.
- But the problem is, if $k > 0$ in Ciucu's result, then our graph G has no matchings.
- So, we modify our region suitably and try to use condensation results.

Our Ideas

- We consider regions with defects (one unit square removed) on the boundaries of an Aztec Rectangle.
- Mills-Robbins-Rumsey's result is for defects on one of the longer sides.
- It is well known, that domino tilings of Aztec Diamonds/Rectangles correspond to counting the number of perfect matchings of its dual planar graph.
- So, from now on we use the terms matchings and tilings equivalently.
- We would like to use Ciucu's result to give a Pfaffian for the regions we are interested in.
- But the problem is, if $k > 0$ in Ciucu's result, then our graph G has no matchings.
- So, we modify our region suitably and try to use condensation results.

But first we look at the case when $k = 0$.

$k = 0$: Aztec Diamond

Here G is now the dual graph of Aztec Diamond and a_i 's are the defects on the boundary.

$k = 0$: Aztec Diamond

Here G is now the dual graph of Aztec Diamond and a_i 's are the defects on the boundary. We need to compute $M(G - \{a_i, a_j\})$.

$k = 0$: Aztec Diamond

Here G is now the dual graph of Aztec Diamond and a_i 's are the defects on the boundary. We need to compute $M(G - \{a_i, a_j\})$.

- If a_i, a_j are on the same side, then the defects are of same type and we get no matchings.

$k = 0$: Aztec Diamond

Here G is now the dual graph of Aztec Diamond and a_i 's are the defects on the boundary. We need to compute $M(G - \{a_i, a_j\})$.

- If a_i, a_j are on the same side, then the defects are of same type and we get no matchings.
- If a_i, a_j are on opposite sides, then the defects are of same type and we get no matchings.

$k = 0$: Aztec Diamond

Here G is now the dual graph of Aztec Diamond and a_i 's are the defects on the boundary. We need to compute $M(G - \{a_i, a_j\})$.

- If a_i, a_j are on the same side, then the defects are of same type and we get no matchings.
- If a_i, a_j are on opposite sides, then the defects are of same type and we get no matchings.
- If a_i, a_j are on adjacent sides, then the defects are of different type and we get matchings. This is given in the next result.

Proposition

Let a, i, j be positive integers such that $1 \leq i, j \leq a$, then the number of domino tilings of $AD(a)$ with one defect on the southeastern side at the i -th position counted from the south corner and one defect on the northeastern side on the j -th position counted from the north corner as shown in the next figure is given by

$$2^{a(a-1)/2} \binom{a-1}{i-1} \binom{a-1}{j-1} {}_3F_2 \left[\begin{matrix} 1, 1-i, 1-j \\ 1-a, 1-a \end{matrix} ; 2 \right].$$

Aztec Diamond with defects on adjacent sides




Figure 3: Aztec Diamond with defects on adjacent sides; here $a = 6$, $i = 4$, $j = 4$

We use Kuo condensation, with the vertices marked as follows.

We use Kuo condensation, with the vertices marked as follows.




Figure 4: Aztec Diamond with some marked squares; here $a = 6$

We use induction with respect to a . The base case of induction is $a = 2$. We would also need to check for $i = 1, j = 1, i = a$ and $j = a$ separately.

We use induction with respect to a . The base case of induction is $a = 2$. We would also need to check for $i = 1, j = 1, i = a$ and $j = a$ separately. If $a = 2$, then the only possibilities are $i = 1$ or $i = a$ and $j = 1$ or $j = a$, so we do not have to consider this case, once we consider the other mentioned cases.

We use induction with respect to a . The base case of induction is $a = 2$. We would also need to check for $i = 1, j = 1, i = a$ and $j = a$ separately. If $a = 2$, then the only possibilities are $i = 1$ or $i = a$ and $j = 1$ or $j = a$, so we do not have to consider this case, once we consider the other mentioned cases.

We now note that when either i or j is 1 or a , some dominoes are forced in any tiling and hence we are reduced to an Aztec rectangle of size $(a - 1) \times a$. It is easy to see that our formula is correct for this.

In the rest of the proof we assume $a \geq 3$ and $1 < i, j < a$. Let us now denote the region we are interested in this proposition as $AD_a(i, j)$. Using the dual graph of this region and applying Kuo condensation with the vertices as labelled in the previous figure we obtain the following identity (details in the next figure),

In the rest of the proof we assume $a \geq 3$ and $1 < i, j < a$. Let us now denote the region we are interested in this proposition as $AD_a(i, j)$. Using the dual graph of this region and applying Kuo condensation with the vertices as labelled in the previous figure we obtain the following identity (details in the next figure),

$$\begin{aligned} M(AD_a(i, j)) M(AD(a-1)) &= M(AD(a)) M(AD_{a-1}(i-1, j-1)) \quad (3.1) \\ &\quad + M(\mathcal{AR}_{a-1, a}(j)) M(\mathcal{AR}_{a-1, a}(i)). \end{aligned}$$

Forced dominoes for different choices of labels




Figure 5: Forced dominoes, where the vertices we remove are marked

Forced dominoes for different choices of labels




Figure 6: Forced dominoes, where the vertices we remove are marked

Simplifying the previous equation, we get the following

$$M(\text{AD}_a(i, j)) = 2^a M(\text{AD}_{a-1}(i-1, j-1)) + 2^{a(a-1)/2} \binom{a-1}{j-1} \binom{a-1}{i-1} \quad (3.2)$$

Simplifying the previous equation, we get the following

$$M(\text{AD}_a(i, j)) = 2^a M(\text{AD}_{a-1}(i-1, j-1)) + 2^{a(a-1)/2} \binom{a-1}{j-1} \binom{a-1}{i-1} \quad (3.2)$$

Now, we use our inductive hypothesis on this equation and we are done.

Simplifying the previous equation, we get the following

$$M(\text{AD}_a(i, j)) = 2^a M(\text{AD}_{a-1}(i-1, j-1)) + 2^{a(a-1)/2} \binom{a-1}{j-1} \binom{a-1}{i-1} \quad (3.2)$$

Now, we use our inductive hypothesis on this equation and we are done.

We now look at the general case.

$k > 0$: Aztec Rectangles

- In order to create a region that can be tiled by dominoes we have to remove k more white squares than black squares along the boundary of $\mathcal{AR}_{a,b}$.

$k > 0$: Aztec Rectangles

- In order to create a region that can be tiled by dominoes we have to remove k more white squares than black squares along the boundary of $\mathcal{AR}_{a,b}$.
- There are $2b$ white squares and $2a$ black squares on the boundary of $\mathcal{AR}_{a,b}$. We choose $n + k$ of the white squares that share an edge with the boundary and denote them by $\beta_1, \beta_2, \dots, \beta_{n+k}$ (we will refer to them as defects of type β).

$k > 0$: Aztec Rectangles

- In order to create a region that can be tiled by dominoes we have to remove k more white squares than black squares along the boundary of $\mathcal{AR}_{a,b}$.
- There are $2b$ white squares and $2a$ black squares on the boundary of $\mathcal{AR}_{a,b}$. We choose $n + k$ of the white squares that share an edge with the boundary and denote them by $\beta_1, \beta_2, \dots, \beta_{n+k}$ (we will refer to them as defects of type β).
- We choose any n squares from the black squares which share an edge with the boundary and denote them by $\alpha_1, \alpha_2, \dots, \alpha_n$ (we refer to them as defects of type α).

$k > 0$: Aztec Rectangles

- In order to create a region that can be tiled by dominoes we have to remove k more white squares than black squares along the boundary of $\mathcal{AR}_{a,b}$.
- There are $2b$ white squares and $2a$ black squares on the boundary of $\mathcal{AR}_{a,b}$. We choose $n + k$ of the white squares that share an edge with the boundary and denote them by $\beta_1, \beta_2, \dots, \beta_{n+k}$ (we will refer to them as defects of type β).
- We choose any n squares from the black squares which share an edge with the boundary and denote them by $\alpha_1, \alpha_2, \dots, \alpha_n$ (we refer to them as defects of type α).
- We consider regions of the type $\mathcal{AR}_{a,b} \setminus \{\beta_1, \dots, \beta_{n+k}, \alpha_1, \dots, \alpha_n\}$, which are more general than the type considered by other authors.

Preliminaries

We define the region $\mathcal{AR}_{a,b}^k$ to be the region obtained from $\mathcal{AR}_{a,b}$ by adding a string of k unit squares along the boundary of the southeastern side (γ defects) as shown in the figure below.

Preliminaries

We define the region $\mathcal{AR}_{a,b}^k$ to be the region obtained from $\mathcal{AR}_{a,b}$ by adding a string of k unit squares along the boundary of the southeastern side (γ defects) as shown in the figure below.




Figure 7: $\mathcal{AR}_{a,b}^k$ with $a = 4, b = 8, k = 4$

Theorem

Assume that one of the two sides on which defects of type α can occur does not actually have any defects on it. We assume this to be the southwestern side.

Theorem

Assume that one of the two sides on which defects of type α can occur does not actually have any defects on it. We assume this to be the southwestern side. Let $\delta_1, \dots, \delta_{2n+2k}$ be the elements of the set $\{\beta_1, \dots, \beta_{n+k}\} \cup \{\alpha_1, \dots, \alpha_n\} \cup \{\gamma_1, \dots, \gamma_k\}$ listed in a cyclic order.

Theorem

Assume that one of the two sides on which defects of type α can occur does not actually have any defects on it. We assume this to be the southwestern side. Let $\delta_1, \dots, \delta_{2n+2k}$ be the elements of the set $\{\beta_1, \dots, \beta_{n+k}\} \cup \{\alpha_1, \dots, \alpha_n\} \cup \{\gamma_1, \dots, \gamma_k\}$ listed in a cyclic order.

Then we have

$$M(\mathcal{AR}_{a,b} \setminus \{\beta_1, \dots, \beta_{n+k}, \alpha_1, \dots, \alpha_n\}) = \frac{1}{[M(\mathcal{AR}_{a,b}^k)]^{n-k+1}} \text{Pf}[(M(\mathcal{AR}_{a,b}^k \setminus \{\delta_i, \delta_j\}))_{1 \leq i < j \leq 2n+2k}],$$

where all the terms on the right hand side are given by explicit formulas.

We need to show that each quantity on the right hand side can be computed.

We need to show that each quantity on the right hand side can be computed.

- $M(\mathcal{AR}_{a,b}^k \setminus \{\beta_i, \beta_j\}) = 0,$

We need to show that each quantity on the right hand side can be computed.

- $M(\mathcal{AR}_{a,b}^k \setminus \{\beta_i, \beta_j\}) = 0,$
- $M(\mathcal{AR}_{a,b}^k \setminus \{\alpha_i, \alpha_j\}) = 0,$

We need to show that each quantity on the right hand side can be computed.

- $M(\mathcal{AR}_{a,b}^k \setminus \{\beta_i, \beta_j\}) = 0,$
- $M(\mathcal{AR}_{a,b}^k \setminus \{\alpha_i, \alpha_j\}) = 0,$
- $M(\mathcal{AR}_{a,b}^k \setminus \{\gamma_i, \gamma_j\}) = 0,$

We need to show that each quantity on the right hand side can be computed.

- $M(\mathcal{AR}_{a,b}^k \setminus \{\beta_i, \beta_j\}) = 0,$
- $M(\mathcal{AR}_{a,b}^k \setminus \{\alpha_i, \alpha_j\}) = 0,$
- $M(\mathcal{AR}_{a,b}^k \setminus \{\gamma_i, \gamma_j\}) = 0,$
- $M(\mathcal{AR}_{a,b}^k \setminus \{\alpha_i, \gamma_j\}) = 0,$

We need to show that each quantity on the right hand side can be computed.

- $M(\mathcal{AR}_{a,b}^k \setminus \{\beta_i, \beta_j\}) = 0,$
- $M(\mathcal{AR}_{a,b}^k \setminus \{\alpha_i, \alpha_j\}) = 0,$
- $M(\mathcal{AR}_{a,b}^k \setminus \{\gamma_i, \gamma_j\}) = 0,$
- $M(\mathcal{AR}_{a,b}^k \setminus \{\alpha_i, \gamma_j\}) = 0,$
- $M(\mathcal{AR}_{a,b}^k)$ is given by Aztec Diamond Theorem.




Figure 8: Removing the forced dominoes from $\mathcal{AR}_{a,b}^k$; here $a = 5, b = 10, k = 5$

- It is given by the previous proposition (Aztec Diamond with defects on adjacent sides) if β_i is on the south-eastern side and not above a γ defect;

$$M(\mathcal{AR}_{a,b}^k \setminus \{\beta_i, \alpha_j\})$$

- It is given by the previous proposition (Aztec Diamond with defects on adjacent sides) if β_i is on the south-eastern side and not above a γ defect;
- Otherwise it is 0,





Figure 9: Choices of β -defects that lead to no tiling of $\mathcal{AR}_{a,b}^k$

$$M(\mathcal{AR}_{a,b}^k \setminus \{\beta_i, \alpha_j\})$$



$$M(\mathcal{AR}_{a,b}^k \setminus \{\beta_i, \gamma_j\})$$

- It is given by Aztec Diamond Theorem if β_i is above a γ defect;

- It is given by Aztec Diamond Theorem if β_i is above a γ defect;
- It is given by the next proposition if the β -defect is in the northwestern side and its distance from the western corner is more than the distance of the γ -defect from the southern corner

Proposition

Let $1 \leq a, i \leq b$ be positive integers with $k = b - a > 0$ and $y = \min\{i, k\}$, then the number of domino tilings of $\mathcal{AR}_{a,b}$ with $k - 1$ squares added to the southeastern side starting at the second position (and not at the bottom) as shown in the next figure and a defect on the northwestern side at the i -th position counted from the western corner is given by

$$2^{a(a+1)/2} \binom{a}{i-y} \sum_{l=0}^{y-1} \binom{a+y-l-2}{y-l-1} {}_3F_2 \left[\begin{matrix} 1, 1-y+l, i-a-y \\ i-y+1, 2-a-y+l \end{matrix}; -1 \right]. \quad (3.3)$$

Regions with defects contd.




Figure 11: $\mathcal{AR}_{a,b}$ with $k-1$ squares added on the southeastern side and a defect on the i -th position shaded in black; here $a=4, b=10, k=6, i=7$

- It is given by Aztec Diamond Theorem if β_i is above a γ defect;
- It is given by the previous proposition if the β defect is in the northwestern side at a distance of more than $k - 1$ from the western corner;

- It is given by Aztec Diamond Theorem if β_i is above a γ defect;
- It is given by the previous proposition if the β defect is in the northwestern side at a distance of more than $k - 1$ from the western corner;
- It is given by the next proposition if the β defect is on the southeastern side;

- It is given by Aztec Diamond Theorem if β_i is above a γ defect;
- It is given by the previous proposition if the β defect is in the northwestern side at a distance of more than $k - 1$ from the western corner;
- It is given by the next proposition if the β defect is on the southeastern side;
- Otherwise it is 0.

Proposition

Let $1 \leq a \leq b$ be positive integers with $k = b - a > 0$, then the number of domino tilings of $\mathcal{AR}_{a,b}(j)$ with $k - 1$ squares added to the southeastern side starting at the second position (and not at the bottom) as shown in the Figure 12 is given by

$$2^{a(a+1)/2} \binom{a+k-1}{j-1} \binom{j-2}{k-1} {}_3F_2 \left[\begin{matrix} 1, 1-j, 1-k \\ 2-j, 1-a-k \end{matrix} ; 1 \right]. \quad (3.4)$$

Regions with defects contd.




Figure 12: Aztec rectangle with $k - 1$ squares added on the southeastern side and a defect on the j -th position shaded in black; here $a = 4, b = 10, k = 6, j = 8$

- The proofs of the previous stated propositions, also uses Kuo condensation in various cases.

- The proofs of the previous stated propositions, also uses Kuo condensation in various cases.
- For the most general case, we have a nested Pfaffian form for the number of matchings.

Theorem

Let $\beta_1, \dots, \beta_{n+k}$ be arbitrary defects of type β and $\alpha_1, \dots, \alpha_n$ be arbitrary defects of type α along the boundary of $\mathcal{AR}_{a,b}$. Then $M(\mathcal{AR}_{a,b} \setminus \{\beta_1, \dots, \beta_{n+k}, \alpha_1, \dots, \alpha_n\})$ is equal to the Pfaffian of a $2n \times 2n$ matrix whose entries are Pfaffians of $(2k+2) \times (2k+2)$ matrices of the type in the statement of main theorem.

Thank you for your attention.